


12/28/10

DOC ID 525257

New York State Department  
of Environmental Conservation  
Division of Solid & Hazardous Materials  
Pesticide Product Registration



**EcoPCO® WP•X**  
**WETTABLE POWDER INSECTICIDE**

**ACTIVE INGREDIENTS:**  
2-Phenethyl Propionate ..... 3.00%  
Thyme Oil ..... 5.00%  
Pyrethrins ..... 0.50%  
**OTHER INGREDIENTS:** ..... 91.50%  
100.0%

Ants, beetles, boxelder bugs, carpenter bees, centipedes, cockroaches, crickets, earwigs, firebrats, fleas, fruitflies, millipedes, phoridflies, pillbugs, sciaridflies, scorpions, silverfish, sowbugs, spiders, termites, wasps  
Adelgids, aphids, elm leaf beetle, fire ants, japanese beetles, lace bugs, leafhoppers, mealybugs, thrips, whiteflies

- **Quick Control**
- **For Use Both Indoors and Outdoors**
- **For Structural and Ornamental Pest Control**

**Fresh Scent**

EPA Reg. No. 67425-25-655  
EPA EST. No. 10208-MO-1

**CAUTION  
KEEP OUT OF REACH  
OF CHILDREN**

See Side/Back Panel for Additional  
Precautions and First Aid

**NET CONTENTS: 1 LB**

**PRETISS  
INCORPORATED**

EcoPCO® WP•X is intended for use as a general surface, spot, or crack and crevice application in and around Homes, Offices, Hotels, Motels, Schools, Hospitals, Nursing Homes, Day-care Centers, Factories, Warehouses, Stores, Supermarkets, Restaurants, Food Handling Establishments, Commercial or Institutional Kitchens, Food Processing Plants, Zoos and Transportation Equipment (Aircraft (do not use in aircraft cabin), Buses, Trucks, Trailers, Boats, Ships, and Trains.) EcoPCO WP•X is intended to be mixed with water and applied with a hand-pressurized or power-operated sprayer. Spray pressure should not exceed 50 psi at the nozzle tip when used indoors.

EcoPCO® WP•X will not stain or cause damage to any painted or varnished surface, plastic, fabric, or other surface when water applied alone causes no damage.

**DIRECTIONS FOR USE:**

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

**Application Equipment Preparation:** For best results, thoroughly wash out sprayer and screens with water and detergent before using EcoPCO WP•X. To avoid sprayer clogging, do not use strainers or screens finer than 50 mesh size. Mixing Procedures: Carefully open container by holding it near its base, avoid squeezing the container as this causes contents to puff-out at the user.

Quantities to be used are given in volumetric terms (tablespoons or cups) and also by weight (grams and ounces). The product is supplied with a measuring scoop (1 cup).

Using the scoop included with the package, scoop out the desired amount of product for each application. When finished, rinse the scoop with water, rinsate should go in tank. Dry and return scoop to the container. Close container.

**For best results, use EcoPCO WP•X as follows:**

1. Fill applicator tank 1/2 with water.
2. Add desired EcoPCO WP•X.
3. Allow 1 to 2 minutes for powder to wet.
4. Agitate, stir, or shake applicator tank.
5. Finish filling applicator tank and agitate again before use to ensure satisfactory suspension.

Note: Upon standing without agitation for more than 1 hour, the applicator tank should be agitated, stirred, or shaken again to achieve satisfactory suspension. Make up only amount of application volume as required. This product should not be mixed with oil.

**CRAWLING AND FLYING PESTS APPLICATIONS**

Crawling & Flying Pests	Concentration of Active Ingredient	Dilution rate
Ants, Boxelder Bugs, Centipedes, Cockroaches (maintenance), Crickets, Earwigs, Firebrats, Fleas, Millipedes, Pillbugs, Silverfish, Sowbugs, Termites**	0.0125%	3 tablespoons or 15 grams (0.5 oz) per gallon of water

Crawling & Flying Pests	Concentration of Active Ingredient	Dilution rate
Ants, Bees, Carpenter Ants, Carpenter Bees, Cockroaches (clean-out), Fire ants (mounds)*, Scorpions, Spiders, Wasps	0.05%	12 tablespoons, 3/4 of a cup or 60 grams (2 oz) per gallon of water

\* For outdoors use only\*\* Above ground only

**Boxelder Bugs, Cockroaches, Sowbugs, Pillbugs, Spiders, Crickets, Scorpions, Millipedes, Centipedes, Silverfish, and Firebrats:** Apply as a coarse, low-pressure treatment to areas where these pests hide or where they may enter the premises. Applications should include cracks/crevices and void areas around and behind interior and exterior walls, cabinets, baseboards, moldings, sinks, pipe openings, storage areas, closets, and appliances. Repeat as necessary. Also see "Outdoor Surfaces Use".

**Ants and Carpenter Ants:** Apply as a coarse, low-pressure treatment to any trails, around doors and windows, around and behind cabinets, walls and similar areas where ants may be found. Pay particular attention to cracks/crevices and void areas where ants can enter the premises. Apply perimeter barrier treatment to prevent infestation as described below. Repeat as necessary. Also see "Outdoor Surfaces Use".

**Bees, Carpenter Bees and Wasps:** Thoroughly treat nest and entrance, surfaces, and surrounding areas where these insects alight. May be used inside residential buildings as well as in and around carports, garages and storage buildings. Also see "Outdoor Surfaces Use".

**Fire Ants (mounds):** Apply as a drench. Gently sprinkle (watering can) a total of 1 gallon over the surface of mounds up to 1 inch in diameter and 2 gallons for larger mounds. Thoroughly wet mound and surrounding areas. Treat new mounds as they appear. All colonies in the vicinity should be treated.

**Scorpions:** Treat and remove accumulations of lumber, firewood, and other materials that serve as harborage sites. Apply as a coarse spray to the point of drip. Also see "Outdoor Surfaces Use".

**Termites Above Ground (Excluding Formosan):** Apply to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is vulnerable. Applications may be made to inaccessible areas by drilling and then injecting with a crack and crevice injector into the damaged wood or void spaces. Applications to attics, crawl spaces, unfinished basements, or man-made voids may be made with a coarse fan spray to control workers and winged reproductive forms of termites in mud shelter tubes. This type of application is not intended to be a substitute for soil treatment. It is intended to kill localized infestations of termites.

**Food-Handling Establishments:** Places other than private residences in which food is held, processed, prepared, or served. Non-Food Areas: EcoPCO WP•X may be used as a general, spot or crack and crevice treatment in non-food areas. Food Areas: EcoPCO WP•X is limited to crack and crevice and spot treatments in food areas. Exposed food should be covered or removed. Do not apply directly to food or food-handling surfaces.

**STORED PRODUCT AND PANTRY PESTS APPLICATIONS**

Stored Product and Pantry Pests	Concentration of Active Ingredient	Dilution rate
Beetles (Cadelle, Cigarette, Confused flour, Dermistid, Drug-store, Granary weevil, Hide, Larder, Leather, Lesser grain borer, Lesser mealworm, Merchant grain, Mealworm, Red flour, Rice weevil, Sawtoothed grain, Warehouse moths, Indian meal, Mediterranean flour), Fruitflies, Phoridflies, Sciaridflies	0.0125%	3 tablespoons or 15 grams (0.5 oz) per gallon of water

**Pantry Pests (of exposed adult and immature stages of insects):** Make application to cupboards, shelving and storage areas. Particular attention should be given to treating cracks and crevices and other harborage areas. Remove all utensils, uncovered foodstuffs (or any with package opened), shelf paper and other objects before treating. Allow treated surfaces to dry and cover shelves with clean paper before replacing utensils, foodstuffs or other items.

**Stored Product Pests (of exposed adult and immature stages of insect pests):** Apply to cracks, crevices and other surfaces where the pests have been seen or have harborage. Treat warehouses, production facilities, storage areas, rail cars, truck beds and other areas where products are stored before filling with the product. EcoPCO WP•X may be used by pest control operators and grain producers to treat grain storage facilities and other areas noted above for stored product pest control. Cleaning of all areas prior to treatment will increase levels of control.

**OUTDOOR SURFACES USE**

For control of ants, bees, centipedes, cockroaches, crickets, earwigs, firebrats, fleas, flies, millipedes, pillbugs, scorpions, silverfish, sowbugs, spiders, wasps and other similar perimeter insect pests. Apply with either a hand or power application equipment as a residual treatment to surfaces of buildings, porches, screens, door and window frames, eaves, patios, garages and other similar areas where these pests are found. Also, may be used for

applications to landscaped areas in and around residential, commercial, institutional, and industrial buildings where these pests are active.

**Barrier Treatment:** To help prevent infestation of buildings, apply to a band of soil and vegetation 6 to 10 feet wide around and adjacent to the building. Also, treat the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Apply as a coarse spray treatment to thoroughly and uniformly wet the foundation and/or band area, using 1 to 5 gallons of treatment solution applied to 800 to 1600 square feet. Use 1.5 to 3 cups of EcoPCO WP•X per gallon of water. (Concentration of spray solution 0.1 to 0.2% pyrethrin)

#### GENERAL INFORMATION FOR LIVESTOCK PEST CONTROL

**Livestock Housing Structures (including Poultry Houses) and Pet Kennels:** Apply as a general surface and/or crack and crevice spray. Control will be enhanced when facilities are cleaned and interior applications are supplemented with exterior perimeter treatments. Also see "Outdoor Surfaces Use". Do not make applications to animal feedstuffs or watering equipment.

#### GENERAL INFORMATION FOR TURF AND LANDSCAPE ORNAMENTAL PEST CONTROL

Crop	Pest	Dilution rate
Trees, Shrubs, Evergreens, Flowers, Foliage plants, Ground covers	Adelgids, Aphids, Elm leaf beetle, Japanese beetles, Lace bugs, Leafhoppers, Mealybugs, Sawfly larvae, Thrips, White flies	3 tablespoons or 15 grams (0.5 oz) per gallon of water

#### ORNAMENTAL APPLICATIONS

**Foliar Applications:** Start treatments prior to establishment of high pest populations and reapply on an as needed basis.

**Broadcast applications:** Mix required amount of product in sufficient water to uniformly and accurately cover the area being treated. Do not use less than 2 gallons of water per 1000 sq. ft. For optimum control, irrigate thoroughly to incorporate EcoPCO WP•X insecticide into the upper soil profile.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling.

##### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment wash water or rinsate.

##### FIRST AID

**IF SWALLOWED:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Have a person sip a glass of water if able to swallow. Do not give anything by mouth to an unconscious person. **IF INHALED:** Remove person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **IF ON SKIN OR CLOTHING:** Take off the contaminated clothing. Rinse the skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice

##### HOT LINE NUMBER:

Have the product container or label with you when calling a poison control center or doctor, or when going for treatment. You may also contact 1-800-222-1222 (Poison Control Center) for emergency medical treatment information.

#### STORAGE AND DISPOSAL

**Pesticide Storage/Disposal:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. In case of a spill sweep up material and dispose of on site or at a waste disposal facility.

**Container Disposal:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

#### WARRANTY & DISCLAIMER

**Limitation of Liability:** To the extent consistent with applicable law, Prentiss Incorporated makes no warranties of merchantability, or of fitness for a particular purpose, nor any other express or implied warranty except as stated above.

Sold by  
Prentiss, Incorporated  
3600 Mansell Road, Suite 350, Alpharetta, GA 30022  
All rights reserved. Made In USA.

Made in USA  
U.S. Patent No. 5,439,690 / 5,693,344 / 6,004,569  
U.S. and Foreign Pat. Pending  
CODE PCW10XXZ (USA)

Revised 02182010

EcoPCO® is a trademark of EcoSMART Technologies, Inc.