

ACCEPTED
VIA NOTIFICATION
LABEL NOT REVIEWED

AUG 31 2010

New York State Department
of Environmental Conservation
Division of Solid & Hazardous Materials
Pesticide Product Registration

Doc ID: 523851

Prescription Treatment® brand

2% Propoxur Bait

KILLS: Cockroaches, Crickets and Ants (excluding Carpenter, Fire, Harvester and Pharaoh)

ONLY FOR USE IN AND AROUND: Apartment Buildings, Government Offices, Homes, Institutions, Public Buildings, and Non-food/feed Areas* of Food Processing, Poultry Packing and Red Meat Packing Plants, Restaurants, Stores and Warehouses

ACTIVE INGREDIENTS:

Propoxur 2.0%

OTHER INGREDIENTS: 98.0%

TOTAL: 100.00%



**KEEP OUT OF THE REACH OF CHILDREN
CAUTION**



FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 min. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15 - 20 min. Remove contact lenses, if present, after the first 5 min, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

Have a product container or label with you when calling a poison control center or doctor, or going for treatment. You may also call 1-800-832-HELP for emergency medical treatment information.

NOTE TO PHYSICIAN: Product contains a cholinesterase inhibitor. If symptoms of cholinesterase inhibition are present, atropine sulfate in large therapeutic doses is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. **SYMPTOMS OF POISONING:** A sense of "tightness" in the chest, sweating, contracted pupils, stomach pains, vomiting and diarrhea.

See back panel for Directions for Use and Storage and Disposal.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Avoid breathing dust.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and other handlers must wear long-sleeved shirt and long pants and shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife and aquatic invertebrates. Birds and other small mammals feeding on treated bait may be killed. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

EPA Reg. No. 499-518 • EPA Est. No. 7969-MO-1

NET WEIGHT: 4 lbs

81040384



100721-20

2% Propoxur Bait

DIRECTIONS FOR USE **IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT** **IN A MANNER INCONSISTENT WITH ITS LABELING**

A ready-to-use product; kills ants (excluding Carpenter, Fire, Harvester and Pharaoh) (except in California), crickets and the cockroaches listed below. Crickets and roaches die after feeding on the bait. Apply as needed to control the target pests.

- Cockroaches: American, Brown, Brown banded, German and Oriental roaches
- Crickets
- Ants (excluding Carpenter, Fire, Harvester and Pharaoh) (except in California)

USE RESTRICTIONS

- Do not formulate this product into other end use products.
- Do not apply to floors or other surfaces likely to be contacted by children or pets.
- Do not apply this product in a way that will contact any person or pet, either directly or indirectly. Keep people and pets out of the application area during application.
- Do not apply as a landscape treatment (to lawns, shrubs or trees, garden plants).

INDOOR USE

For use in Apartment Buildings, Government Offices, Homes, Institutions, Public Buildings, Restaurants (non-food areas), Stores and Warehouses.

Apply bait lightly to floors near baseboards, in closets, under sinks and refrigerators, around and inside garbage cans, and in other areas where insects are found but which are inaccessible to children such as cracks and crevices. (Do not make Crack & Crevice® applications in California.) Do not apply to animals. Do not treat portions of buildings where animals can feed on bait. Use approximately 4 oz of bait per 1,000 ft² of space and scatter lightly so that bait is barely visible. Repeat as necessary. Keep out of reach of children.

FOOD/FEED HANDLING ESTABLISHMENTS (Non-food Areas Only)

Food/feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served.

*Non-food/feed areas, where applications of this product are permissible, are garbage rooms, entries and vestibules, lavatories, floor drains, offices, locker rooms, boiler rooms, machine rooms, garages, mop closets, storage after canning/bottling, warehouses where food/feed is not exposed, buses, boats, ships, trains and cargo areas of planes.

Apply bait on paper, pasteboard, or other material that will permit removal. Locate these makeshift bait stations on window sills, on floor near walls, in storage areas and other areas where insects have been observed. Maintain strict account of location and number of bait stations so they can be retreated or removed in the future.

RESEARCH FACILITIES

The use of this product in research animal quarters for pest control will not induce changes in liver microsomal enzyme activity in these animals. Do not apply bait directly on animals nor where animals can feed on the bait.

OUTDOOR USE

Use only around patios, driveways, sidewalks and foundations. Apply bait evenly in a band 2 - 3 ft wide at the rate of 4 oz/1,000 ft² around foundations, patios, driveways, or sidewalks, taking care to treat any cracks or crevices. Also treat beneath concrete drain splash pans.

STORAGE AND DISPOSAL

Do not puncture or incinerate! Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed below. In spill or leak incidents, keep unauthorized people away.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Empty container by using the product in accordance with the label directions. Offer empty container for recycling, if available, or place in trash or sanitary landfill or incinerate by burning, if allowed by state and local authorities/regulations. If burned, stay out of smoke. If container is partly filled, contact your local solid waste agency for disposal instructions.

Manufactured for:
Whitmire Micro-Gen Research Laboratories Inc.®
by BASF Corporation
3568 Tree Court Industrial Blvd.
St. Louis MO 63122-6682
Questions? Call 1-800-777-8570
© 2008 Whitmire Micro-Gen Research Laboratories, Inc.®