

# ULD® BP-300

## Contact Insecticide II

For indoor or outdoor application as a direct, space, area or contact spray. Dependent upon the insects and areas to be treated, this product may be applied through mechanical aerosol (non-metered), Ultra Low Volume (ULV) generators, as well as thermal and conventional fogging or spraying equipment.

**KILLS:** Ants (excluding Harvester & Pharaoh), Bed Bugs, Cockroaches, Fleas, Flies, Scorpions, Spiders, Stored Product Pests (Beetles, Mites, Moths) and Ticks

**FOR USE IN:** Commercial (including Food Handling Establishments, Food Service, Storage, Preparation and Processing Areas), Institutional and Residential Areas

See page 2 for additional insects and use sites.

### ACTIVE INGREDIENTS:

Pyrethrins\*, a botanical insecticide ..... 3.0%  
Piperonyl butoxide\*\* ..... 15.0%

### OTHER INGREDIENTS:

Contains Petroleum Distillate ..... 82.0%  
TOTAL: 100.0%

\* 1 gal contains 0.211 lb of pyrethrins and 1.053 lb piperonyl butoxide

\*\* (butylcarbityl) (6-propylpiperonyl) ether and related compounds

EPA Reg. No. 499-522

KEEP OUT OF REACH OF CHILDREN

## CAUTION

### FIRST AID

**IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15 – 20 min. Remove contact lenses, if present, after the first 5 min, then continue rinsing eyes. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 min. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this product (including health concerns, medical emergencies or pesticide incidents), you may call 1-800-832-HELP.

**NOTE TO PHYSICIAN:** May pose an aspiration pneumonia hazard. Contains petroleum distillate.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye irritation. Do not get in eyes, on skin or clothing. Wear protective eyewear\*. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

\*Protective eyewear includes goggles, face shield and/or safety glasses.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):** Mixers, loaders, applicators and other handlers must wear the following: Long-sleeve shirt, long pants, shoes and socks, and chemical-resistant gloves made of barrier laminate, nitrile rubber, neoprene rubber or Viton.

• In addition to the above PPE, applicators using a high pressure hand wand in an enclosed area must wear at least a NIOSH-approved respirator with: a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any R, P, or HE filter.

• In addition to the above PPE, applicators fogging this product (applying as a space spray) in an enclosed area must wear a half-face, full-face, or hood-style NIOSH-approved respirator with: a dust/mist filtering cartridge (MSHA/NIOSH approval number prefix TC-21C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a cartridge or canister with any R, P or HE filter.

**USER SAFETY REQUIREMENTS:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

### USER SAFETY RECOMMENDATIONS

#### User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

**TERRESTRIAL APPLICATIONS:** This product is toxic to aquatic organisms, including fish and invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product has a potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

Except as specified in the directions for use, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

### PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame. Do not apply this product around electrical equipment due to the possibility of shock hazard.

### DIRECTIONS FOR USE

**IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.**

#### USE RESTRICTIONS

- Apply this product only as specified on this label.
- Do not apply more than 1 time/day.
- Only protected handlers may be in the treatment area during application.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Do not remain in treated area. Exit area immediately and do not enter or allow others to enter treated area until sprays have dried, until vapors, mists and aerosols have dispersed, and the treated area has been thoroughly ventilated.
- Remove pets, birds and cover fish aquariums before spraying.
- Do not make outdoor applications during rain.
- Do not use in outdoor residential misting systems.
- Except when applying to sewers or drains as instructed on this label, do not apply directly to sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.
- Do not use in aircraft cabins.
- Remove or cover exposed food and drinking water, in the treatment area, before application.
- Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces, in the treatment area, or wash them before use.
- Do not apply as a space spray in residences.
- For space spray applications in food handling and processing facilities:
  - Do not make space spray applications when facility is in operation.
  - Cover or remove food, in the treatment area, prior to space spray applications.
  - Cover food processing surfaces, in the treatment area, or clean after treatment and before use.

- When used in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- Do not apply this product in hospital rooms while occupied or in any rooms occupied by the elderly or infirm. Remove patients from room prior to treatment and thoroughly ventilate (2 hr) after application. Do not return patients to room until after ventilation.
- Do not use in patient rooms in hospitals and nursing homes for treatment of bed bugs.
- Do not apply to classrooms when in use.

### PRODUCT INFORMATION

**Note:** Some furniture finishes, particularly plastics, can be damaged by petroleum based insecticides. To avoid damage, cover those items and do not spray directly. If in doubt, test by applying this material to an inconspicuous area before spraying.

### USE IN FOOD HANDLING ESTABLISHMENTS

Applications are permitted in food/feed areas of food/feed handling establishments as a contact spray, space spray or Crack & Crevice® treatment, including those operating under the Federal meat, poultry, shell egg grading and egg products inspection programs.

Food handling establishments are places other than private residences in which food is held, processed, prepared or served.

**FOOD AREAS:** Include areas for receiving, serving, storing (dry, cold, frozen, raw), packing (canning, bottling, wrapping, boxing), preparing (cleaning, slicing, cooking, grinding), edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups).

**NON-FOOD AREAS:** Include areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling).

Apply according to specific directions for pest(s) to be treated under the Indoor Use section below. Do not make space spray applications to food areas when the facility is in operation. Prior to space spray applications, remove or cover exposed food and drinking water in the treatment area. Remove or cover dishes, utensils, food processing equipment, and food preparation surfaces in the treatment area, or wash them before use.

### INDOOR USE

**SPACE SPRAY –FLYING INSECTS (Flies & Gnats):** Use full strength or dilute as desired using the dilution chart below. Prior to treatment, open cabinets, doors and equipment to be treated; turn off air conditioners and fans, extinguish all flames and close windows in the area to be treated. Apply using a mechanical aerosol generator (no droplets > 50 microns in diameter and 80% < 30 microns). Apply undiluted at a rate of approx. 1 fl oz/1,000 ft<sup>3</sup> of room space, or apply diluted at the rate of 1 - 2 fl oz/1,000 ft<sup>3</sup> of room space. In commercial operations, subtract the space occupied by stored product or equipment from the cubic footage. Disperse toward areas suspected of harboring the greatest infestations of listed pests. Disperse in all locations (including around equipment, appliances and pallets, if desired) contacting as many insects as possible. Do not remain in treated areas after application. Keep area closed for at least 1 hr. Open and thoroughly ventilate treated areas before reoccupying. Do not apply as a space spray in residences.

**NOTE:** This specimen label is for informational purposes only. All uses may not be approved in all states. See labeling which accompanied product for Directions for Use or call 800-777-8570 for more information.

# ULD® BP-300 Contact Insecticide II

**DILUTION:** Use a white mineral oil conforming to regulation 21 CFR 178.3620(b), or an odorless light petroleum hydrocarbon conforming to 21 CFR 172.882 or 40 CFR 180.910 (inert ingredients used pre and post-harvest; exemptions from the requirement of a tolerance), and 180.930 (inert ingredients applied to animals; exemptions from the requirement of a tolerance). Combine oil and concentrate as directed below and mix well just prior to application.

Pyrethrins % Desired	Mixture Rate	Application Rate/1,000 ft <sup>3</sup>
1.00	1 Part Conc. to 2 Parts Oil	1 fl oz
0.50	1 Part Conc. to 5 Parts Oil	1 fl oz
0.30	1 Part Conc. to 9 Parts Oil	1 - 2 fl oz
0.15	1 Part Conc. to 19 Parts Oil	1 - 2 fl oz

**SPACE SPRAY –CRAWLING INSECTS (Ants (excluding Harvester & Pharaoh), Bed Bugs, Beetles, Earwigs, Cockroaches, Silverfish, Spiders):** Use full strength. Prior to treatment, open cabinets, doors and equipment to be treated; turn off air conditioners and fans, extinguish all flames and close windows in the area to be treated. Apply as a space spray using a mechanical aerosol generator (no droplets >50 microns in diameter and 80% <30 microns) at a rate of approx. 1 fl oz/1,000 ft<sup>3</sup> of room space. In commercial operations, subtract the space occupied by stored product or equipment from the cubic footage. Disperse toward areas suspected of harboring the greatest infestations of listed pests. Disperse in all locations (including around equipment, appliances and pallets, if desired) contacting as many insects as possible. Do not apply as a space spray in residences.

Do not remain in treated area after application and leave area closed for 1 hr. Ventilate thoroughly before occupants are allowed to reenter.

**DIRECT SPRAY OR VOID TREATMENTS:** Use equipment designed for space or mist treatments. Follow the manufacturer's recommendations for proper equipment set up and operation. Place the tip of the applicator at or into the crack or crevice or opening to the void. Apply in short bursts into area being treated. Apply 1 sec of product for every 3 ft<sup>3</sup> of void area. **For Direct Spray,** apply the product in short bursts into the area where insects are to be contacted, such as corners of the room, behind and under furniture or other equipment. Apply from a distance of 18 - 24" away from the surface and allow the material to drift to area to be treated. The amount of product applied is not to exceed 0.125 fl oz/2 ft<sup>2</sup>. **For Bed Bugs:** Direct application into areas where bed bugs harbor. Do not direct this product at mattresses.

**HANDTANK OR PINSTREAM APPLICATION:** Direct application into cracks and crevices in elements of construction, walls, floors, underneath sinks, behind pipes and other places where insects harbor. Spot treatment is also permitted. For spot treatments, use a coarse low-pressure spray. Limit individual spot treatments to an area no larger than 20% of the surface area. Any individual spot treatment shall not exceed 2 ft<sup>2</sup>. Contact as many insects as possible. Ensure product will not stain surface or leave an unacceptable residue prior to application. The amount of product applied is not to exceed 0.125 fl oz/2 ft<sup>2</sup>.

## OUTDOOR USE

**OUTDOOR GROUND FOGGING APPLICATION USING HAND HELD EQUIPMENT (Flies & Gnats):** Use in open areas near buildings and in campgrounds. Use preferably at a time when temperature is cool (≤ 75°F) and the wind velocity is approx. ≤ 5 mph. When using ULV equipment, apply at the rate of 4 fl oz/acre. Allow spray drift to penetrate dense foliage.

## FOR USE ON DOMESTIC ANIMALS/LIVESTOCK

Use to protect beef cattle, dairy cattle and horses from horn flies, house flies and gnats, apply a light mist sufficient to wet the tips of the hair. To kill stable flies on beef cattle, dairy cattle and horses, apply 2 fl oz/adult animal, sufficient to wet the hair but not to soak the hide. To kill the mottled stages of blood sucking lice on beef and dairy cattle, apply the spray to thoroughly wet the hair. Use 2 fl oz or less/animal and repeat every 2 - 3 weeks as required, but do not apply more than 1 time/day. **Do not use on dogs and cats.**

**Space Treatment Over Animals in Barns and Corrals or in Indoor Misting Systems Used in Commercial Barns, Stables and Animal Quarters:** Select proper dilution and treat with ULD equipment over animals, or apply through Indoor Misting Systems.

## USE RESTRICTIONS FOR INDOOR MISTING SYSTEMS

- Not for use in outdoor residential misting systems (indoor or outdoor).
- Do not apply this product in barns or stables where animals intended for slaughter or human consumption will be maintained.
- Do not apply when food, feed and/or water is present.
- When using this product, installers and service technicians must comply with the license, certification or registration requirements of the state(s), tribe(s) or local authority(ies) where they are installed.
- When applying via a remote activation device, do not apply when people and pets are present. If possible, when applying via automatic timer, set the timing for application when people and pets are unlikely to be present.
- Direct nozzles to spray towards the target area and away from areas where people are typically present.
- Do not use in evaporative cooling system.
- Do not use in misters located within 3 ft of air vents, air conditioner units or windows.
- If used in a system with a reservoir tank for the end use dilution, the system reservoir tank must be locked. Securely attach the end use pesticide label and a dilution statement to the system reservoir tank in a weather protected area or plastic sleeve. The dilution statement must be phrased as follows: this container holds \_\_\_ parts [product name] to \_\_\_ parts water.

- If used in a direct injection system, the pesticide container must be locked. Securely attach the end use label to the pesticide container in a weather protected area or plastic sleeve.
- This product must only be used in systems that have been calibrated to apply no more than the maximum application rate of 0.0033 lbs/1,000ft<sup>3</sup> (no animals present) [diluted to 1% = 6.5 fl oz/1,000 ft<sup>3</sup> = 0.0703 lbs/gal] or 0.0008 lbs/1,000 ft<sup>3</sup> (animals present) [diluted to 1% = 0.0005 lbs/fl oz = 0.0703 lbs/gal].

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in secured, dry storage area. Do not store above 100°F for extended periods of time. Storage below 32°F may cause freezing. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Empty container by using the product according to the label directions. Do not reuse or refill this container! Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

Empty the remaining contents into application equipment or a mix tank and drain for 10 sec after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 sec. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 sec after the flow begins to drip. Repeat this procedure 2 more times.

Then offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned stay out of smoke.

**In addition to previously listed sites, this product is acceptable for use in and around the following sites:**

- |                               |                                 |
|-------------------------------|---------------------------------|
| <b>INDOOR:</b>                | Peanut Storage Warehouses       |
| Animal Areas & Quarters       | Poultry Houses                  |
| Auditoriums                   | Poultry Processing Plants       |
| Bakeries                      | Prisons                         |
| Beverage Plants               | Public Buildings                |
| Boats                         | Rabbit Processing Plants        |
| Bottling Plants               | Rabbit Houses                   |
| Brandy Storage Warehouses     | Railroad Cars                   |
| Buildings                     | Recreational Vehicles           |
| Buses                         | Research Animal Areas           |
| Cabins                        | Restaurants                     |
| Canneries                     | Restrooms                       |
| Cattle Barns                  | Rice Mills                      |
| Churches                      | Shade Houses                    |
| Closets                       | Ships                           |
| Condominiums                  | Ship holds                      |
| Communication Centers         | Stables                         |
| Correctional Facilities       | Stable Bins                     |
| Dairy & Livestock Barns       | Stored Seed Warehouses          |
| Dried Fruit Processing Plants | Stores                          |
| Drains & Sewers               | Supermarkets                    |
| Dried Fruit Warehouses        | Swine Houses                    |
| Drugstores                    | Textile Mills & Warehouses      |
| Dumpsters                     | Theaters                        |
| Dwellings                     | Tobacco Plants                  |
| Egg Processing Plants         | Tobacco Warehouses              |
| Factories                     | Townhouses                      |
| Flour Mills                   | Trains                          |
| Food Handling Establishments  | Trucks                          |
| Food Processing Plants        | Truck Trailers                  |
| Freight Containers            | USDA Inspected Facilities       |
| Fruit Packing Sheds           | Utilities                       |
| Garbage Cans                  | Utility Rooms                   |
| Garbage Compactors            | Voids (Attics, Walls, Ceilings) |
| Grain Elevators               | Warehouses                      |
| Grain Mills                   | Wineries                        |
| Grain Handling Equipment      | Wine Storage Warehouses         |
| Granaries                     | Zoos                            |
| Homes                         | <b>OUTDOOR:</b>                 |
| Horse Stables/Barns           | Campgrounds                     |
| Hospices                      | Corrals                         |
| Industrial Installations      | Drive-in Restaurants            |
| Institutions                  | Drive-in Theaters               |
| Jails                         | Dumpsters                       |
| Laboratories                  | Feedlots                        |
| Libraries                     | Garbage Dumps                   |
| Livestock Barns               | Golf Courses                    |
| Loafing Sheds                 | Junkyards                       |
| Locker Rooms                  | Kennels                         |
| Mills                         | Landscapes                      |
| Mausoleums                    | Lath Houses                     |
| Meat Packing Plants           | Manure Piles                    |
| Milk Rooms                    | Parks                           |
| Missions                      | Plantscapes                     |
| Mobile Homes                  | Playgrounds                     |
| Museums                       | Recreational Areas              |
| Mushroom Processing Plants    | Sewers                          |
| Nursing Homes                 | Swineryards                     |
| Offices                       | USDA Inspected Facilities       |
| Office Buildings              | Zoos                            |
| Passenger Rail Cars           |                                 |
| Peanut Processing Plant       |                                 |

**This product is also effective against the following insects when used according to label directions:**

## FLYING INSECTS:

Bees  
Blow Flies  
Bottle Flies  
Cheese Skippers  
Deer Flies  
Flies  
Fungus Gnats  
House Flies  
Hornets  
Mushroom Flies  
Stable Flies  
Vinegar Flies  
Yellowjackets

## GRAIN BORERS:

Larger Grain Borers  
Lesser Grain Borers

## CRAWLING INSECTS:

Bed Bugs  
Book Lice  
Cabbage Loopers  
Cabbage Worms  
Cockroaches  
Crickets  
Fire Ant Workers  
Firebrats  
Millipedes  
Pillbugs  
Scorpions  
Sowbugs

## FLEAS & TICKS:

Brown Dog Ticks  
Fleas  
Ticks  
**SPIDERS AND MITES:**

Cheese Mites  
Clover Mites  
Grain Mites  
Mites  
Spiders

## LICE:

Cattle Lice

## WORMS:

Dark Mealworms  
Lesser Mealworms  
Yellow Mealworms

## BEETLES:

Black Carpet Beetles  
Carpet Beetles  
Darkling Beetles  
Dermestid Beetles  
Dried Fruit Beetles  
Drugstore Beetles  
Flat Grain Beetles

## FOREIGN GRAIN BEETLES:

Foreign Grain Beetles  
Fungus Beetles  
Hede Beetles  
Ground Beetles  
Merchants Grain Beetles  
Mexican Grain Beetles  
Red Flour Beetles  
Red Horned Grain Beetles  
Rusty Grain Beetles  
Sawtoothed Grain Beetles  
Spider Beetles  
Trogoderma Beetles  
White Marked Spider Beetles

## WEEVILS:

Bean Weevils  
Granary Weevils  
Maize Weevils  
Rice Weevils

## MOTHS:

Almond Moths  
Chocolate Moths  
Clothes Moths  
Cocoa Bean Moths  
Mediterranean Flour Moths

## CONDITIONS OF SALE AND WARRANTY

Follow the **Directions for Use**. It is impossible to eliminate all risks inherently associated with use of this product, and therefore all such risk shall be assumed by the Buyer. Whitmire warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions for Use**, subject to the inherent risks, referred to above. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW: (A) WHITMIRE MAKES NO OTHER WARRANTIES EXPRESS OR IMPLIED, INCLUDING WARRANTIES OF FITNESS FOR PARTICULAR PURPOSE OR MERCHANTABILITY, (B) BUYER'S EXCLUSIVE REMEDY AND WHITMIRE'S AND SELLER'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT, AND (C) WHITMIRE AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, INCIDENTAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. Whitmire and the Seller offer this product, and the Buyer accepts it, subject to these **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of Whitmire.

Manufactured by:  
Whitmire Micro-Gen Research Laboratories, Inc.®  
by BASF Corporation  
3568 Tree Court Industrial Blvd.  
St. Louis MO 63122-6682  
Questions? Call 1-800-777-8570  
© 2012 Whitmire Micro-Gen Research Laboratories, Inc.®

ULD® and Crack & Crevice® are registered trademarks of BASF Corporation.