PAGE 5 PAGE 4

along baseboards and door and window frames; behind and beneath sinks, stoves, refrigerators and cabinets and around Spray areas where these pests are found or may hide, including dark corners of rooms and closets; cracks and crevices in walls; Psocids, Scorpions, Silverfish, Sowbugs, Spiders, and Waterbugs: To Kill Ants, Centipedes, Cockroaches, Crickets, Earwigs, Millipedes, mit humans or pets to contact treated surfaces until the spray has dried. Do not treat establishments where livestock feed is present. GENERAL: Apply as a coarse wet spray to thoroughly treat hiding places, such as cracks, crevices, and moist areas around pipes. Under refrigerators and along baseboards to control peats listed below. Use at the rate of one (1) gallon of spray mixture per 1,000 aguare feet. Spray directly on insects when possible, Do not per-mix humans and a manufactured and a manufactured

Application Within Residential and Non-Residential Buildings for Application Within Residential and Non-Residential Buildings from of Industrial, Institutional, and Commercial Buildings, such Cars, Hospitals, Hotels, Warehouses, Truck Trailers, Railroad Cars, Theatures, Office Buildings, and Schools. FOR PREMISE PESTS:

pound and rinsed with potable water before using. Do not apply when food processing facility is in operation or foods see a workers or the thersils that sense to surfaces or undersils that may come in contact with food. Food contact surfaces and equipmenty contact should be thoroughly cleaned with an effective cleaning comment should be thoroughly cleaned with an effective cleaning comment and many come.

Соскгоасрея. Saw-Toothed Grain Beetles, Angoumois Grain Moth, Mediterranean Flour Moth, Indian Meal Moth, Cigarette Beetle, Grain Mites and \*Rice Weevil, Yellow Meal Worm, Cadelle, Confused Flour Beetle,

in treated area and ventilate thoroughly before re-entry. Do not repeat application more than once every three (3) days. with other products approved for this use. Use as above in mechanical aerosol foggers at the rate of one (1) ounce per 1,000 cubic feet of space. Keep area closed for at least 1/2 hour. Do not remain Warehouses, Storage Rooms and Similar Locations: Infested stored products should be removed and destroyed, fumigated or treated To Kill The Adult and Accessible Forms of Insect Pests\* Of Stored Food (Stored In Multi-Walled Paper Bags) In

Do not repeat application more than once every three (3) days. 7,000 cubic feet of space. Do not remain in treated area. Keep area closed for at least 1/2 hour and ventilate thoroughly before re-entry.

uers of the room at the rate of one (1) ounce (diluted spray) per ances, cupboards, or any dark harborage areas, then treat as a space spray directing the sprey towards the ceiling and upper corproduces particles of aerosol size thirty (30 microns, or less). Direct spray first into all cracks and crevices, behind all equipment, applicentrate diluted in one (1) gallon of oil only to make a 0.25% spray. Use diluted spray in a mechanical aerosol fogger or generator which room and shut off all air conditioning or ventilating equipment. For space spray applications use a maximum of ten (10) ounces of con-As A Space Spray: To Kill Houseflies, Mosquitoes, Gnats and Small Flying Moths: Close

tourteen (14) days in all types of food handling establishments. than one application every seven (7) days in restaurants and similar food service establishments or more than one application every Use a dosage of one (1) gallon of spray per 1,000 square feet. Contact as many insects as possible with direct spray. Do not exceed more into all cracks and crevices in woodwork, floors, underneath sinks, areas behind pipes and in all places where these insects shelter. Peocids, Scorpions, Silverfish, Sowbugs, Spiders and Waterbugs: Apply this spraywith a sprayer that produces coarse, wetting droplets To Kill Ants, Centipedes, Cockroaches, Crickets, Earwigs, Millipedes,

Do Not Exceed More Than One Application Every Fourteen (14) food; excessive residues in food may result.

facility is in operation or foods are exposed. Do not apply this spray to surfaces or utensils that may come into contact with and into floor drains, non-food conveyors, and other areas where granary insects may be found. Do not apply when food processing feet. Pay particular attention to cracks, crevices and similar pro-tected locations in floors and walls. Spray thoroughly around spray to surfaces where these pests are normally found, such as floors, baseboards, around walls, benches and pieces of equipment. Use at a dosage of one (1) gallon of spray per 1,000 square As a Surface Spray.

To Kill the Accessible Stages of Granary Insects: Rice Weevils,
Contused Flour Beetles, Must Red Flour Beetles, Saw-Tonhed Grain
Beetles, Cadelles, Meal Moth Larvae and Adults, Cigarette Beetles,
Drug Store Beetles, and Granary Weevils: Apply as a coarse wet
Drug Store Beetles, and Granary Weevils: Apply as a coarse wet
somewing the stage of the

after canning or bottling.

drains to sewers, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage

but avoid soaking young or flowering plants. Repeat treatments may be necessary at 7 to 14 day intervals. Kills the <u>lxodes dammini</u> tick (deer tick), which transmits lyme disease. Also kills foraging fire ants upon direct contact. A registered fire ant mound treatment should also be used in conjunction with this product.

Keep pets and children out of the treated area until after the spray has dried. Do not spray animals. Dogs, cats, their kennel or bedding area should also be treated with a registered flea control product in conjunction with this application.

Amount of Product to Use Fluid Ounces	Amount of Water to Use	Area to be Treated
1/6 Fluid Ounce	3 gallons	500 sq. ft. (50' X 10')
1/3 Fluid Ounce	6 gallons	1,000 sq. ft. (50' X 20')
2/3 Fluid Ounce	12 gallons	2,000 sq. ft. (50' X 40')

### OUTDOOR CONTROL OF ASIAN COCKROACHES:

To control outdoor infestations of Asian cockroaches: Dilute 10 teaspoons of Speckoz® Evercide® Esfenvalerate EC in 30 gallons of Asian cockroaches: Dilute 10 teaspoons of Speckoz® Evercide® Esfenvalerate EC in 30 gallons of the control of the co water to treat 5,000 sq. ft. Repeat as needed. Keep childre off treated areas for 24 hours or until the grass has completely dried.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. STORAGE: Store in a warm, dry place. Keep container closed.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. In case of spill or leak, soak up with sand, earth or synthetic absorbents. Do not use alkaline absorbents or clean area of spill with alkaline detergents.

**DISPOSAL:** Do not reuse empty container. Wrap container in several layers of newspaper and discard in trash.

Distributed by: SPECKOZ, INC. 100 North Point Center, Suite 330 Alpharetta, GA 30022

 $\mathsf{EVERCIDE}^{\circledcirc}$  is a Reg. TM of McLaughlin Gormley King Co.  $\mathsf{SPECKOZ}^{\circledcirc}$  is a Reg. TM of Speckoz, Inc.

EPA Reg. No. 1021-1641-72113

EPA Est. No. 1021-MN-2

# tenvalerate EC

### ater or Oil Dilutable

CONTROLS NUMEROUS PESTS IN AND AROUND STRUCTURES RECOMMENDED FOR COMMERCIAL AND INDUSTRIAL USE For Use in and Around the Home and Food and Non-Food Areas of: Schools, Warehouses, Office Buildings, Apartment Buildings, Theatres, Hotels, Motels, Kennels, Food Processing Plants, Food Service Establishments, Truck Trailers, Railroad Cars, and Food Manufacturing and Warehousing Establishments, Also for Use on Lawns to Kill Fleas and Ticks.

ACTIVE INGRÉDIENT:

\* Contains petroleum distillate EPA Reg. No. 1021-1641-72113 Rev. 1204-0406 EPA Est. No. 1021-MN-2

KEEP OUT OF REACH OF CHILDREN CAUTION

0,1011011		
FIRST AID		
	Immediately call a poison control center or doctor. On not induce vomiting unless told to do so by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.	
IF ON SKIN OR CLOTHING:	Take off contaminated clothing.     Rinse skin immediately with plenty of water for 15-20 minutes.     Call a poison control center or doctor for treatment advice.	
IF INHALED:	Move person to fresh air.     If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.     Call a poison control center or doctor for further treatment advice.	
NOTE TO DIMOIOLAN O		

IOTE TO PHYSICIAN: Contains petroleum distillate –vomiting may cause spiration pneumonia. Have the product container or label with you when calling poison control center or doctor, or going for treatment. For information regarding nedical emergencies or pesticide incidents, call 1-888-740-8712.

802662-016

NET CONTENTS 1 PT.

PAGE 8

E JDAY PAGE 6

ing rooms, but excluding areas where foods may be prepared or held. Non-food areas include garbage rooms, lavatories, floor age and enclosed processing systems, mills and dairies. Serving areas are places where prepared foods are served, such as din-Food areas include areas for receiving, storage, packing, canning, bottling, wrapping, boxing, preparing, edible waste stor-

livestock feed is present. Food Manufacturing and Processing Establishments, and Food Warehousing Establishments .Do not treat establishments where and Handling, Burl/Oot Limited to: Restaurants, Grocery Stores, Bakeries, Including, Burl/Oot Limited to: Restaurants, Grocery Stores, Bakeries, Application Only Within Food and Mon-Food Areas of Food Service

of contact sprays use at the rate of one (1) gallon (diluted spray) per 1,000 square feet of surface area. trol or light infestations use one (1) ounce of concentrate per gal-lon. When using oil as the diluent, an odorless light pertoleum hydrocarbon as defined by CBF Title 21, Sec. 172.884 should be used for all tood area and food related applications. For all peptications of contact reserve use a The rate of the III and it contacts and and a contact of the contact o ingredient. For initial clean out of heavy infestations use two (2) our est of concentrate per gallon of spray; for maintenance concentrating the per gallon of spray; for maintenance concentrating the concentration of spray in the concentration of the concentra FOR INDOOR PESTS:

GENERAL: Use Speckoz® Evercide® Estenvalerate EC to control the peockoz® Evercide® Estenvalerate EC to control the pests indicated in the listed areas as a contact spray wixing either one (1) or two (2) fluid ounces of concentrate in one (1) allow on (1) or two (2) fluid ounces of concentrate in one (1) allow of 10 nor one of 10 no

SHAKE WELL BEFORE USE where occupants are present.

has dried.

Do not apply this product in occupied patient rooms or in any rooms/areas while occupied by a patients, the elderly or infirm. Do not apply to classrooms while in use. Do not apply in institutions (including libraries, sports facilities, etc.) in the immediate area whose occurrents are suppresent in the immediate area.

Do not allow children or pets to contact treated surfaces until spray Remove pets, birds, and cover fish aquariums before spraying.

used for food or feed. sils should be covered during treatment or thoroughly washed before use following treatment. Cover exposed food. Do not spray plants

ract ants, cockroaches, crickets, sowbugs, spiders and ticks. Use two (2) fluid ounces per one (1) gallon of water to control on con-FOR OUTDOOR PESTS: Gvercide® Exfenvalerate EC at the rate of

Do not reapply more than every fourteen (14) days. sure to push insecticide deep into the space. Avoid applying to the point of runoff or drip. Ventilate area thoroughly before re-entry. cations. Follow the injection equipment manufacturer's recommendations for the proper set up and air presenes. Place the applicator tip at or into the void space to be treated. For insceessible voids, it may be necessary to drill an access hole(s). Inject product into the void space in a flort bursts, allowing the set presented to the void space in short bursts, allowing the all presented productions in the void space in a flort bursts, allowing the all presented productions. To Treat Voids in Equipment and Structures: To kill Ants (including Carpenter Ants), Centipedes, Cockroaches, Crickets, Earwigs, Psocids, Scorpions, Silverfish, Spiders, Waterbugs and Insect Pests of Stored Food: To kill insects harboring in wall voids and other inaccessible spaces in equipment as attuctures use injection equipment designed for deal void ULU (Illus Low Volume) appliations and estimate Commentation and the commentation of the holes into wall voids where these ants or their nests are present other areas where these ants enter the premises or hide. Spray into cracks and crevices, through openings or small newly drilled To Kill Carpenter Ants: Apply two (2) ounces per one (1) gallon of water as a coarse spray, around doors, door frames, windows and

registered for use on animals. from the area. Do not permit humans or pets to contact treated euror fores until the spray has dried. Old bedding should be removed and replaced with clean, fresh bedding after treatment. Do Not Surey Yers With This Proluce. Treat dogs and cats with a product renistered for use on animals. various Associated or integration of the paterial of the paterial of the control of Theorems or using the one (1) ounce per gallon of water dilution to infested rugs; can per resting areas. Prior to treatment, aduations and fish bould be covered and birds, such as cenarias, removed firsh bounds should be covered and birds. behind baseboards, window and door frames and localized areas of floor and floor coverings where these peats may be present. For thoroughly apply as a spot treatment to infested areas such as pet beds and resting quarters, nearby cracks and crevices, along and To Kill Brown Dog Ticks and Fleas: For Control of Brown Dog Ticks,

garbage cans, plumbing and other utility installations. Pests driven out of hiding pleces should be sprayed directly. For Ants, apply to antitual doors and windows and wherever these pests may find entrance.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **CAUTION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid raminum is wainower, limitated or absorbed indugitismit. Avoid contact with skin, eyes or clothing. Avoid breathing vapors or spray mist. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS
This product is toxic to fish, birds and other wildlife. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not con-taminate ornamental fish ponds. Do not contaminate water when taminate ornamental fish ponds. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product into sewer systems without previously notifying the sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA. PHYSICAL OR CHEMICAL HAZARDS.

## PHYSICAL OR CHEMICAL HAZARDS

**COMBUSTIBLE.** Do not use or store near heat or open flame. Do not apply this product around electrical equipment due to the possibility of shock hazard.

**DIRECTIONS FOR USE**It is a violation of Federal Law to use this product

in a manner inconsistent with its labeling.

Speckoz® Evercide® Esfenvalerate EC can be mixed with an EPA registered insect growth regulator or pyrethrin-containing products in areas where these products are allowed to be sprayed. Please check specific label directions of these products before using. Do not tank mix this product with dichlorvos-(DDVP)-containing products

Do not apply directly to food. Cover or remove all food and food processing equipment during application. After spraying in meat packing plants, bakeries, and other food processing plants, wash all equipment, benches, shelving, etc. where exposed food will be handled with an effective cleaning compound, and then rinse with potable water. In the home, all food processing surfaces and uten-

at the rate of one (1) gallon per 1,000 square feet to outside surfaces of buildings, porches, patios, garages and other areas where these pests have been seen or are found. Do not spray in or near fish ponds or other bodies of water. Avoid soaking young or flowering plants. Building Perimeter Treatment: To help prevent infestations of buildings by ants, crickets, spiders, centipedes, and sowbugs (pillbugs) treat a 12-18 inch band of soil or other substrate adjacent to buildings and the building foundation to a height of 2 to 3 feet where these pests are active and may find entrance. Use two (2) fluid ounces of concentrate per gallon of water and apply as a coarse wet spray at the rate of one (1) gallon per 1,000 square feet.

TO KILL EXPOSED/SWARMING TERMITES: OUTDOORS ONLY. Use two (2) fluid ounces of Concentrate per gallon of water and apply as a coarse, wetting spray when swarming termites are seen emerging from wood piles, wooden fence posts, wooden structures, or from the ground. Swarming usually occurs in the spring or at other times when a termite colony becomes overcrowded and new repro-ductive termites with wings emerge and fly away to mate and establish new colonies. This treatment will control the sprayed termites and will not protect the structure from which the swarm is coming. Use only as a contact spray to kill emerging reproductive (winged) and worker termites emerging from infested wood.

TO KILL WOOD DESTROYING PESTS: OUTDOORS ONLY. Carpenter Bees, Carpenter Ants, Old House Borers, Powder Post Beetles, Furniture Beetles, Round-Headed House Borers, Deathwatch Beetles. Use two (2) fluid ounces of Concentrate per gallon of water and apply as a coarse, wetting spray to exposed pests and to the damaged areas of wood, spraying into galleries or tunnels in the exposed wood. Also spray around doors, door frames, windows and other areas where these pests may enter the house or hide. Spray into cracks and crevices and, if necessary, drill small holes and spray into inacks. cessible wooden structural voids where these pests are suspected LAWN FLEA AND TICK CONTROL: To Kill Fleas and Ticks Harboring in Lawns and Backyards: For power sprayers, dilute and apply as shown in the table below. [Use this concentrate in a hose-end sprayer. Place required amount of concentrate in sprayer jar (see chartbelow) and fill with water to the recommended level. Mix well. Attach to sprayer and spray as recommended.] For best results, lawn should be mowed a day or two before spraying. Spray using a slow, even sweeping motion, being sure to cover the entire lawn surface where the pets will normally frequent. Spray under ornamentals and trees,