



# SAGA® WP INSECTICIDE

Specimen Label

- \* FOR RESIDUAL CONTROL OF MAJOR NUISANCE PESTS
- \* ONLY FOR SALE, USE AND STORAGE BY PEST CONTROL OPERATORS
- \* DESIGNED FOR USE IN FOOD/FEED AND NON-FOOD/NON-FEED AREAS OF FOOD/FEED PROCESSING PLANTS.
- \* FOR USE IN USDA INSPECTED FACILITIES

**ACTIVE INGREDIENT:**

(1R, 3S) 3[(1'RS) (1', 2', 2',2' -tetrabromoethyl)] -2, 2-dimethyl-cyclopropanecarboxylic acid (S)- alpha-cyano-3-phenoxybenzyl ester<sup>1/</sup> . . . . 40.00%

**INERT INGREDIENTS:** . . . . . 60.00%  
100.00%

<sup>1/</sup> Tralomethrin.

PRECAUCION AL CONSUMIDOR: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read English, do not use this product until the label has been fully explained to you.)

EPA REG. NO. 432-755

EPA EST. NO.:

## KEEP OUT OF REACH OF CHILDREN WARNING FIRST AID

- If Swallowed:**
- ◆ Call a poison control center or doctor immediately for treatment advice.
  - ◆ Have person sip a glass of water if able to swallow.
  - ◆ Do not induce vomiting unless told to by a poison control center or doctor.
  - ◆ Do not give anything my mouth to an unconscious person.

- If in Eyes:**
- ◆ Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lens, if present, after the first 5 minutes, then continue rinsing.
  - ◆ Call a poison control center or doctor for treatment advice.

- If on Skin or Clothing:**
- ◆ Take off contaminated clothing.
  - ◆ Rinse skin immediately with plenty of water for 15-20 minutes.
  - ◆ Call a poison control center or doctor for treatment advice.

- If Inhaled:**
- ◆ Move person to fresh air.
  - ◆ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.

Have container label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment information.

For Product Information Call Toll-Free: 1-800-331-2867

See Side Panel For Additional Precautionary Statements

NET CONTENTS:

BAYER ENVIRONMENTAL SCIENCE  
A Business Group of Bayer CropScience LP  
95 Chestnut Ridge Road • Montvale, NJ 07645

## PRECAUTIONARY STATEMENTS

### Hazards To Humans & Domestic Animals

#### WARNING

May be fatal if swallowed. Harmful if absorbed through the skin or inhaled. Causes eye irritation. Avoid breathing dust or spray mist. Avoid contact with skin, eyes or clothing. Contact with product may result in transient tingling and reddening of the skin. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.

#### Environmental Hazards

This pesticide is extremely toxic to fish and other aquatic organisms, including crab and shrimp. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Use with care when applying in areas adjacent to any body of water. Do not contaminate water when disposing of equipment waters.

#### Physical & Chemical Hazards

Do not use water base sprays of SAGA WP in conduits, motor housings, junction boxes, or other electrical equipment because of possible shock hazard.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Mix as needed. Do not store diluted material. Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. If the container is leaking and/or material is spilled, on floor or paved surfaces, sweep up and remove to chemical waste area. Concentrate is stable at normal storage temperatures.

Pesticide Disposal: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Empty containers completely into application equipment. Do not reuse empty container. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### GENERAL INFORMATION:

Saga WP is a wettable powder insecticide containing 40% by weight active ingredient which will provide effective and long residual control of the most pests listed on this label.

Remove pets and cover aquariums and terrariums before spraying.

Saga WP does not damage paints, plastics, fabrics or other surfaces where water alone causes no damage.

Saga WP is intended for use by professional applicators in and around residential and nonresidential structures and their immediate surroundings and on various modes of transportation. Permitted areas of use include, but are not limited to the Food/Feed and Non-Food/Non-Feed Areas of:

aircraft apartment buildings bakeries bottling facilities breweries buses cafeterias candy plants	industrial buildings and installations laboratories marine vessels mausoleums meat, poultry and egg processing and packing plants nursing homes
--	--

canneries food manufacturing, processing and servicing establishments grain mills granaries greenhouses (non-commercial) hospitals hotels houses	railcars restaurants schools ships and vessels stores and institutions trailers trucks warehouses wineries
--	--

Treatments may be repeated as necessary to maintain adequate control.

#### MIXING INSTRUCTIONS:

Saga WP is intended to be mixed with water and applied with hand pressurized or power operated sprayers as a coarse spray. When diluting, add water to the spray tank and the appropriate amount of Saga WP. Thoroughly agitate the sprayer before using to insure proper suspension of the wettable powder. Re- invert sprayer before use if spraying is interrupted. If spray screens are used, they should be 50 mesh or larger. Dilute Saga WP insecticide only with water.

#### DOSAGE CHART:

This product is intended for dilution prior to applications as directed in the table below. Use the scoop provided for measuring from the jar. Each level scoopful contains 0.1 oz. Saga WP.

Pests	Concentration of Active Ingredient	Dilution Rate Per Gallon of Final Spray
ants, centipedes, clothes moths, cockroaches (maintenance rate), crickets, earwigs, flies, firebrats, gnats, midges, fleas (contact), millipedes, mosquitoes, silverfish, spiders	.03% w/w	1 Scoop*
bees, carpet beetles, cockroaches (clean out rate), fleas, hornets, killer bees, pillbugs, scorpions, sowbugs, ticks, yellow jackets, wasps	.06% w/w	2 Scoops
termites (Formosan & Eastern subterranean-localized control only)	.06% w/w	2 Scoops
fire ants, carpenter ants (localized control only)	.06% w/w	2 Scoops

\*1 scoop = 0.1 ounce of Saga WP = 1 level teaspoon.

#### GENERAL PEST CONTROL INDOORS

Treat specific areas that insects inhabit or where they are frequently found. Mix 1 scoop (0.1 ounce) of Saga WP in one gallon of water and apply to adequately cover 1000 square feet. Do not allow dripping or run-off to occur. Under conditions of severe pest infestation, when quick knockdown and/or longer residual control is desired, mix 2 scoops (0.2 ounce) of Saga WP in one gallon of water and apply to adequately cover 1000 square feet. Do not allow dripping or run-off to occur. To insure consistent acceptable control of the target pest, proper sanitation and prevention of reinfestation are recommended.

Note: Do not permit adults, children or pets to contact treated surfaces

until spray has dried completely. Do not apply this product in patient rooms or in any rooms while occupied by the elderly or infirm. Do not apply to classrooms when occupied by students. Do not use as a space spray. Use in ventilated areas. Do not apply where electrical short circuits could occur. Use a dust or a dry bait in these areas.

**PREMISE PEST CONTROL:** *Cockroaches, Ants, Crickets, Silverfish, Firebrats, Centipedes, Millipedes, Clothes Moths, Earwigs, Pillbugs, Sowbugs, Scorpions and Spiders:* Apply as a coarse, low pressure spot or crack and crevice spray to areas that these pests normally inhabit. Pay particular attention to dark corners of rooms and closets; floor drains; cracks and crevices in walls; along and behind baseboards; beneath and behind sinks, stoves, refrigerators, and cabinets; around plumbing and other utility installations; doors, windows and in attics and crawl spaces. Applications may be made to floor surfaces along baseboards and around air ducts.

**PANTRY PEST CONTROL:** To kill adult and accessible stages of insect pests such as *Lesser Grain Borers, Grain Weevils, Flour Beetles, Saw-toothed Grain Beetles, Cigarette Beetles, Merchant Grain Beetles, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, Grain Mites and Spider Mites.*

Make application to cupboards, shelving and storage areas. Remove all food stuffs, all utensils, shelf paper and other objects before spraying. Allow all treated surfaces to dry and cover shelves with clean paper prior to replacing food stuffs, utensils or other items. Any food stuff accidentally contaminated with any spray solution should be discarded.

**ANTS (including Pharaoh Ants):** Apply to trails, around doors and windows and other areas where ants inhabit or enter premises.

**FLEAS:** A total management program is recommended. It is suggested that infested areas be thoroughly vacuumed prior to treatment. These areas include all carpet, upholstered furniture, baseboards and other indoor areas frequented by pets. The vacuum cleaner bag should be discarded in an outdoor trash container. Pet bedding should be cleaned or replaced.

Apply Saga WP to rugs, carpets, upholstered furniture, pet beds, floor coverings, pet resting areas and other infested habitats. The quantity of finished spray required is dependent upon carpet/fabric density. Provide uniform coverage, apply until infested area is moist.

Effective *Flea* control includes controlling the source of flea infestations. This can require treatment of pets with an EPA approved product registered for such use and treating outside areas frequented by pets. Saga WP can also be applied out-of-doors for flea control. Do not treat pets with this product. Do not allow pets to contact treated areas until spray has dried. Remove animal and feed before spraying.

**TICKS (INCLUDING LYME DISEASE VECTORS):** Thoroughly apply Saga WP to infested areas, such as pet beds and resting quarters; nearby cracks and crevices; along baseboards, windows and door frames, where these pests may be present. For best results, old bedding should be replaced with clean bedding after treatment. Do not treat pets with this product. Do not allow pets to contact treated areas until spray has dried. Remove animals and feed before spraying. Saga WP can also be applied out-of-doors for tick control.

**TERMITES, CARPENTER ANTS AND CARPENTER BEES:** (For localized control only). Apply spray dilution to voids or channels in damaged wooden members of a structure, or to cracks, spaces or bearing joints between wooden members of a structure or between such members and the foundations, in locations vulnerable to attack such as crawl spaces. Drilling may be necessary prior to applying Saga WP to voids, cracks or joint spaces in vulnerable locations and to channeled sections of wood but it must be recognized that such drilling and treating of a timber in place may not result in complete coverage of infested and vul-

nerable voids or galleries. All treatment holes drilled in construction elements in commonly occupied areas of structures must be securely plugged. Inspect at 6 months and 1 year to insure that the colony has been killed.

The purpose of such applications of Saga WP is to kill workers or winged reproductive forms which may be present in the treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment but are merely a supplement. The purpose of such applications of Saga WP for termites is to kill workers or reproductive forms of the insect.

Do not apply solution until location of heat or air conditioning ducts, vents, water and sewer lines and electrical conduits are known and identified. When located, applications may be made around these structures. However, caution must be taken to avoid contamination of these structural elements and airways.

**AIRCRAFT PEST CONTROL:** Do not use in aircraft cabins. Apply to cracks, crevices, and other surfaces where the labeled pests have been seen or may have harborage. Allow spray to dry before re-entry.

#### FOOD/FEED HANDLING ESTABLISHMENTS

For control of all labeled pests, applications are permitted in food/feed and non-food/non-feed areas of food/feed handling and processing establishments as a general surface, spot or crack and crevice treatment.

Food/Feed handling establishments are defined as places other than private residences in which exposed food/feed is held, processed, prepared or served. Included also are areas for receiving, storing, packing (canning, bottling, wrapping, boxing), preparing, edible waste storage and enclosed processing systems (mills, dairies, edible oils, syrups) of food/feed. Serving areas where food/feed is exposed and the facility is in operation are also considered food/feed areas.

**NOTE:** Containers larger than one pound may not be marketed or stored in food handling establishments.

#### NON-FOOD/FEED AREAS

Examples of nonfood/nonfeed areas include garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, mop closets and storage (after packaging, canning or bottling). All areas that insects inhabit or through which insects may enter should be treated.

**USDA INSPECTED FACILITIES:** Saga WP is authorized by USDA for use in federally inspected meat and poultry plants. This product is acceptable for use in edible product areas of official establishments operating under the Federal Meat, Poultry, Shell Egg Grading and Egg Products Inspection Programs, as a crack and crevice treatment. Saga WP is classified as an F2 residual insecticide according to regulations listed under Section 5.6(A)(1)(b) of the USDA Food Safety and Inspection Service Handbook. Refer to List of Proprietary Substances and Nonfood Compounds for authorized listing.

**GENERAL SURFACE APPLICATION:** Do not use this application method in the food/feed handling areas of the facility when the facility is in operation or foods are exposed. Do not apply directly to food/feed products. Cover or remove all food/feed processing and/or handling equipment during application. After application in food/feed processing plants, cafeterias and similar facilities, wash all equipment, benches, shelving and other surfaces which food/feed will contact. Clean food/feed handling or processing equipment and thoroughly rinse with clean, fresh water. After application in bakeries, clean all food/feed contact surfaces thoroughly with appropriate equipment.

**SPOT, OR CRACK AND CREVICE APPLICATIONS:** In USDA Inspected facilities, apply only when facility is not in operation. In non-USDA facilities, applications may be made while the facility is in operation provided

exposed food/feed is covered or removed from area being treated prior to application. Do not apply directly to food/feed or food/feed-handling surfaces.

In the home, do not allow spray to contact food/feed surfaces, if it does, clean surfaces with soap and water.

#### GENERAL PEST CONTROL OUTDOORS

Use Saga WP to control outdoor pests by application as either a perimeter treatment or as a residual spray.

**PERIMETER TREATMENTS:** To help prevent infestation of buildings, apply Saga WP where pests are active and may find entrance. For example, treat a band of soil 6 to 10 feet wide around and adjacent to buildings and treat the building foundation to a height of 2 to 3 feet. Apply as a coarse spray at the rate of 1 scoop or 2 scoops of Saga WP in sufficient water to adequately cover 1000 square feet (consult dosage chart). Thoroughly and uniformly wet the barrier area.

**RESIDUAL SPRAY:** To control *Ants, Bees, Cockroaches, Crickets, Hornets, Spiders, Wasps* and *Yellow Jackets (contact kill)*, apply Saga WP as a coarse low pressure spray to surfaces of buildings, porches, screens, window frames, eaves, patios, lawns (residential only), refuse dumps and garages. For application to *Bees, Hornets, Wasps* and *Yellow Jackets* apply late in the evening when insects are at rest.

**TO CONTROL *Fire Ants*** apply as a mound drench at a rate of 2 scoops (0.2 ounce) per gallon of water. Sprinkle (watering can) one gallon diluted product over the surface of each mound up to 12 inches in diameter and two gallons for larger mounds. Thoroughly wet mound and surrounding area. If possible, applications should be made on a warm day after rainfall.

**TO CONTROL *Fleas And Ticks (including Lyme Disease Vectors, Ticks-contact Kill)*** apply Saga WP as a coarse wet spray to areas where fleas and ticks are most likely to be found, such as, but not limited to, dog houses, kennels, runways, cracks and crevices in sidewalks, paths or patios, under plants, shrubs, trees, or other shaded, moist areas where pests may rest.

**FLIES, MIDGES, GNATS, AND MOSQUITOES:** (for use as an aid in reducing annoyance from these insects) Spray outside surfaces of doors, screens, window frames, or wherever these insects may enter the room. Also, spray surfaces around light fixtures, on porches, in garages and other places where these insects alight or congregate.

**CARPENTER ANTS:** To control inside trees, utility poles, fencing and decking materials and similar structural members, drill to find the interior infested cavity and inject 0.06% dilution using appropriate treatment tool with splashback guard. To control tunneling Carpenter Ants in soil, apply 0.06% dilution as a drench or inject at intervals of 8 to 12 inches, using 1 quart per linear foot. Establish a uniform barrier at the edge of walls, driveways, or other hard surfaces where ants are tunneling beneath the surfaces. To protect firewood from carpenter ants and termites, treat soil beneath firewood prior to stacking with a 0.06% dilution at 1 gallon per 8 square feet to prevent infestation.

**ORNAMENTAL PESTS:** To control *Armyworms, Bagworms, Cankerworms, Cutworms, Elm Leaf Beetles, Flea Beetles, Grasshoppers, Gypsy Moth Larvae, Adult Japanese And June Beetles, Lacebugs, Leafhoppers, Leafrollers, Leaf Skeletonizers, Loopers, Oakworms, Plant Bugs, Pine Moths, Sawfly Larvae, Spanworms, Stinkbugs, Tent Caterpillars, Ticks, Treehoppers, Webworms.*

Use Saga WP at 1 scoop per 20 gallons of water (plus spreader/sticker) and spray foliage thoroughly but not the point of run-off.

Not For Use In Commercial Greenhouses Or Nurseries.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on

plants intended for aesthetic purposes or climate modification and being grown in ornamental parks and gardens or lawns and grounds.

**LAWN AND GROUNDS PESTS:** To control *Ants Armyworms, Chinchbugs, Crickets, Cutworms, Sod Webworms, Fleas* and *Ticks.*

Use Saga WP at 1 scoop per 5 gallons of water and spray at 4 gallons per 1000 square feet.

Not for use on sod farms, grass grown for seed or on golf course turf.

#### TANK MIXING (INDOORS AND OUTDOORS)

**GENERAL:** Saga WP can be tank mixed with pesticides currently registered for similar use patterns, unless the companion product label specifically prohibits such mixing. Tank-mix applications must be made in accordance with the more restrictive of label limitations and precautions. No label application rates may be exceeded. Prior to tank-mixing, a small jar compatibility test should be conducted using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture. When tank-mixing with Saga WP, first add the Saga to the diluent and agitate to thoroughly suspend it in solution, then add the prescribed amount of companion product.

In combination with KICKER:

For enhanced flushing and knockdown of *Cockroaches*, tank mix at the rate of 1/4 to 1/2 ounce (equivalent to 1/2 to 1 tablespoon or 7.4 to 14.8 ml) of Kicker per gallon of finished spray and apply as directed above.

For improved control of pyrethroid resistant *German Cockroaches*, tank mix at the rate of 2 to 4 ounces of Kicker per gallon and apply as directed above.

For knockdown of *Adult Fleas*, tank mix at the rate of 0.5 to 1 ounce of Kicker per gallon and apply as directed above.

**IN COMBINATION WITH INSECT GROWTH REGULATORS (IGR'S):** To extend control of immature stages of *Cockroaches* and *Fleas*, tank mix the recommended rate of Saga WP with the correct amount of IGR per finished gallon. Follow all directions for use patterns and spray intervals from the IGR label to achieve maximum benefit.

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond Bayer Environmental Science's control, it is impossible for Bayer Environmental Science to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** BAYER ENVIRONMENTAL SCIENCE MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer Environmental Science is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. Bayer Environmental Science disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** THE EXCLUSIVE REMEDY OF THE USER OR

BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER ENVIRONMENTAL SCIENCE'S ELECTION, THE REPLACEMENT OF PRODUCT.

©Bayer AG, 2003

SAGA is a registered trademark of Bayer AG.

BAYER ENVIRONMENTAL SCIENCE  
A Business Group of Bayer CropScience LP  
95 Chestnut Ridge Road  
Montvale, NJ 07645

PRODUCT OF FRANCE

SWP-SL-01/03